

# **Longitudinal, Base-to-Apex Myocardial Perfusion Abnormalities in patients with coronary artery disease.**

Hiroki Otani, Hideaki Endo, Yutaka Kagaya, Kunio Shirato Kenji Tamaki<sup>1)</sup>, Eiji Nozaki<sup>1)</sup>  
Toshiaki Sasaki<sup>2)</sup>, Kazunori Terasaki<sup>2)</sup>, Tsuneo Takahashi<sup>3)</sup>

Department of Cardiovascular Medicine, Tohoku University Graduate School of Medicine  
2-1 Seiryu Aoba, Sendai, 980-8575, Japan

1) Department of Cardiovascular Medicine, Iwate Prefectural Central Hospital  
1-4-1 Ueda, Morioka, 020-0066, Japan

2) Cyclotron Research Center, Iwate Medical University  
348 Tomegamori, Takizawa, 020-0173, Japan

3) Takizawa Institute, Japan Radioisotope Association  
348 Tomegamori, Takizawa, 020-0173, Japan

## **Abstract**

Gould et al reported that diffuse coronary artery narrowing causes graded, progressively decreasing perfusion along the base-to-apex, longitudinal axis of the heart after dipyridamole. However, frequency and severity of longitudinal perfusion abnormality is unclear in patients with coronary artery disease who undergo coronary arteriography. We performed positron emission tomography with <sup>13</sup>N ammonia in 9 patients with coronary artery disease (angina pectoris =7, myocardial infarction =2). Eight patients underwent coronary angioplasty of diseased vessel. <sup>13</sup>N ammonia PET was performed at rest and after dipyridamole. The infarcted territories were excluded for analysis. All analyzed myocardial territories were perfused by coronary artery with less than 75% luminal narrowing. Longitudinal perfusion abnormalities was greater in myocardial territory which is perfused by angioplasty-performed vessel compared with that perfused by non angioplasty-performed vessel. We need further study to clarify clinical significance of longitudinal myocardial perfusion abnormality in patients with coronary artery disease.