PIXE analysis of mothers' and infants' hairs collected at medical checkups held in Fukuoka city

S. Kuroda³, H. Imaizumi³, S. Goto², T. Yamada⁴, T. Takatsuji³, T. Nakamura³, K. Sera⁵ and Y. Nose¹

¹Graduate School of Health Sciencies, Kumamoto Health Sciencies University 325 Izumi-machi Kumamoto 861-5598 Japan

²Nishina Memorial Cyclotron Center, Japan Radioisotope Association 348-58 Tomegamori, Takizawa, Iwate 020-0173, Japan

³Graduate School of Science and Engineering, Nagasaki University 1-14 Bunkyo-machi, Nagasaki 852-8521, Japan

> ⁴Mie University Graduate School of Medicine 2-174 Edobashi, Tsu, Mie 514-8507, Japan

²Cyclotron Research Center, Iwate Medical University 348-58 Tomegamori, Takizawa, Iwate 020-0173, Japan

Abstract

This is the third report of our project to examine relationships between atopic dermatitis and hair minerals of infants and mothers. The hair minerals are measured using the proton-induced x-ray emission (PIXE) method. The second report published last year described the characteristics the hair minerals of mothers and children and some differences of them between one month and ten months after birth. This study examined the accuracy of the PIXE measurements by statistically analyzing two independent measurements obtained from different hairs of the same person.

The project is expected to proceed as follows:

- $1) \ The \ results \ of \ the \ PIXE \ measurements \ will \ be \ reported \ to \ the \ participation \ doctors.$
- 2) Association analysis of the amount of minerals, dining habit, and clinical performance.
- 3) Statistical analysis for causal relationships between mineral deficiency/excess and atopy/allergy conditions.
- 4) Redefine the participating mother/child population as a new cohort to study the effects of environment, dining and minerals on their health conditions.
- 5) Cooperation with the "Birth cohort project" by the Ministry of Environment.
- 6) Cooperation of the birth cohort study by Norway