

Imaging brain amyloid using the radioligand ^{18}F -AV45 (Florbetapir F18)

T. Shibata¹, H. Yonezawa¹, S. Takahashi¹, J. Takahashi¹, M. Kudoh¹,
S. Obara¹, Y. Terayama¹, T. Sasaki², K. Terasaki² and K. Sera²

¹Department of Neurology and Gerontology, Iwate Medical University
19-1 Uchimaru, Morioka, Iwate 020-8505, Japan

²Cyclotron Research Center, Iwate Medical University
348-58 Tomegamori, Takizawa, Iwate 020-0173, Japan

Abstract

【Purpose】 To characterize quantitative ^{18}F -AV45 (Florbetapir) positron emission tomographic (PET) measurements of fibrillar β -amyloid (A β) burden of participants with probable Alzheimer disease (AD), mild cognitive impairment (MCI), healthy control (HC), cerebral amyloid angiopathy (CAA) and frontotemporal lobar degeneration (FTLD).

【Methods】 ^{18}F -AV45 was performed on 9 patients with AD (men 5, women 4; HDS-R 18 ± 5.8 points, mean age 72y), and 3 patients with MCI (men 2, woman 1; HDS-R 27 ± 2.0 points, mean age 62y), and 3 healthy controls (men 2, woman 1; mean age 70y), 2 patients with CAA (man 1, woman 1), 4 patients with FTLD (man 1, women 3; HDS-R 25 ± 4.3 , mean age 72y). Dynamic PET was performed over approximately 90 min after injection of the tracer (370 MBq). Subsequently, we constructed time-activity curves. Standardized uptake values and cortical-to-cerebellum standardized uptake value ratios (SUVRs) were calculated with regions of interest (ROI) at cortical region (frontal, temporal, parietal, occipital lobe) and other regions (putamen, thalamus, Pons) for all subjects.

【Results】 The standardized uptake value ratio (SUVR) calculated using the cortical-to-cerebellar ratio in patients with AD, MCI and HC was plateaued within 40 min after ^{18}F -AV-45 administration. The cortical SUVR for 10 minutes extending from 50 to 60 minutes after administration was 1.30-1.37 for the patients with AD, 1.12-1.29 for the patients with MCI, 1.03-1.26 for the HC. The SUVR for the patients with AD were greater than those for the HC. SUVR value for the MCI was middle average. The images among AD patients with SUVR value between 0.6 to 2.0 were visually discriminative from HC. SUVR of white matter were higher than that of gray matter for the HC while that of gray matter was higher for the patients of AD.

【Conclusion】 ^{18}F -AV-45 PET showed significant difference between AD patients and HC, utilizing simplified SUVR acquired by scanning of 50-60 minutes after ^{18}F -AV-45 administration.