

Environmental audit of mercury-free operation at artisanal/small-scale mining sites in Mongolia

Satoshi Murao¹, Shoko Goto², Jambardorj Uramгаа³, Baatar Tumenbayar³,
Sainbileg Minjin⁴ and Koichiro Sera⁵

¹National Institute of Advanced Industrial Science and Technology
1-1-1 Higashi, Tsukuba 305-8567, Japan

²Nishina Memorial Cyclotron Center, Japan Radioisotope Association
348-58 Tomegamori, Takizawa, Iwate 020-0603, Japan

³Sans Frontiere Progres
P.O.Box 46, Ulaanbaatar – 46, Mongolia

⁴MICHIKO Foreign Language Training and Translation Center
Ulaanbaatar, Mongolia

⁵Iwate Medical University Cyclotron Research Center
348-58 Tomegamori, Takizawa, Iwate 020-0603, Japan

1 Introduction

In spite of the strict environment policy, mercury contamination is still present in various environmental media and food (especially fish) all over the globe. One of the reason for such contamination is the artisanal/small-scale gold mining (ASGM) where informal gold-ore beneficiation releases mercury due to the amalgamation. According to UNEP¹⁾, mercury release from ASGM is estimated to be about 1,400 t/year making it the largest global demand sector for mercury.

In order to mitigate the use of mercury in the ASGM sector, “ethical jewelry” is recently promoted as an economic incentive to the miners²⁾. The ethical jewelry requires responsible conducts to all of the stakeholders in the supply chain and secures fair payment to the players. Some mining groups began to follow this idea and announced that they do not use mercury any more³⁾.

PIXE at NMCC is, because of the physical features, highly expected to be an ideal tool for the environmental audit to confirm the right conducts at ASGM operations, and thus the authors harnessed the system to verify that a group in Mongolia who assert mercury-free work is obeying the standards and rules.

2 ASGM in Mongolia

In gold producing regions in Mongolia always seen are artisanal/small-scale miners who dig the ground or pan the river sediment (Figs. 1 and 2). They are called *Ninja* after Japanese cartoon *Ninja*

Turtles because they carry plastic water basin on their back like turtle's shell. Such informal mining started around 2000⁴⁾ and is now rampant all over the nation⁵⁾.



Fig. 1 An artisanal underground mining site in Mongolia where they follow ecological gold rules.



Fig. 2 An example of artisanal placer gold mining in Mongolia. Some workers are female.

Recently some miners who learnt eco-friendly way of small-scale mining started to introduce mercury-free technology and disseminate it to other groups. They invite Ninjas who use mercury to consider the environment and human rights and to join their NGO which designates *Artisanal and Small Scale Mining National Federation of Mongolia*. This is a big umbrella which holds many ASGM groups like SAM Project group in this study.



Fig. 3 Office of Artisanal and Small Scale Mining National Federation of Mongolia in Ulaanbaatar



Fig. 4 An example of mercury-free ore dressing plant which was set up by Swiss ODA.

The SAM group is a miners' organization³⁾ which was initially assisted by Swiss Agency for Development and Cooperation (SDC) through its *Sustainable Artisanal Mining (SAM)* project⁶⁾ to get information and mercury-free ore dressing plants (Figs. 3 and 4). They assert that they do not use any toxic chemicals except for small amount of acid to dissolve impurities in gold concentrate.

3 Samples

Environmental samples were obtained in and around the ASGM site in Bayanhongor Aimag

and from an ore dressing plant in Mandal community, Zuunkharaa Soum, Selenge Aimag. In Bayanhongor, a SAM project site was visited and workers' hairs were taken after getting their informed consent. Also Nariniiguur community of Bayan-Ovoo Soum near the SAM site was visited and local people's hairs were obtained after getting informed consent (Appendices I and II). Foodstuff in a Mongolian-style meat soup was also sampled. At Mandal plant (Fig. 4), tailings were taken after getting permission from the person in charge of the plant.

4 Analysis

Food stuff was dried and ground in an agate mortar till mixture is a fine powder. A slight portion of the powder was taken and put on the 4.0 μm polypropylene film and was fixed with 3 μl of 1.0 % collodion solution as a tiny patch.

The targets were bombarded with a 2.9 MeV proton beam extracted from a small-sized cyclotron at NMCC, and the emitted X-rays were simultaneously measured with two Si(Li) detectors. X-ray absorbers were used for the No. 1 detector. 500- μm -thick Mylar film was used for the food, while 300- μm Mylar was used for the hair samples.

As is shown on Table 1 foodstuff showed no contamination of heavy metals. All of the human hairs indicated no mercury anomaly but higher concentration of arsenic and lead. In the tailings mercury was below the detection limit.

5 Concluding remarks

Influence of the toxic elements seems to be minimal in around the ASGM who asserts the compliance of laws and regulations. Because of the following features, PIXE at NMCC is regarded to be a robust instrument to assist the environmental monitoring and audit of the ethical jewelry.

- ✓ Capability to analyze any type/form of specimen
- ✓ Simultaneous measurement of many elements heavier than sodium. Almost equal sensitivity for most elements.
- ✓ Nondestructive analysis
- ✓ Analysis of untreated specimens
- ✓ Short measurement time typically less than five minutes for one specimen

References

- 1) UNEP (2013) Global Mercury Assessment, UNEP Division of Technology, Industry and Economics, Chemicals Branch International Environment House, Geneva, 32pp.
- 2) Murao, S. (2013) Present status of ethical jewelry, *Geo-Pol. Sci. Med. Geol. And urban Geol.* 9, 9-17 (in Japanese with English abstract).
- 3) Murao, S., Tumenbayar, B., Uramгаа, J. and Minjin S. (2015) Present status of ethical gold production in Mongolia, *Proc. 25th Symp. Geo-Environments and Geo-Techniques*, 121-122.
- 4) Tumenbayar, B., Murao, S., Sera, K., Futatsugawa, S., Nukushina, R., Grayson, R. and Maidar, T. (2001) Small-scale mining in Mongolia –a prompt report-, *Chishitsu News No.564*, 49-51 (in Japanese).
- 5) Tumenbayar, B., Murao, S., Maidar, T., Uramгаа, J. and Sera, K. (2013) Anthropogenic mercury contamination and geology of Mongolia, *Proc. 25th Symp. Geo-Environments and Geo-Techniques*, 109-112.
- 6) Khuldorj, B., Murao, S. and Nakano, A. (2015) Lessons learnt from an international cooperation of artisanal gold mining in Mongolia: a case study of Sustainable Artisanal Mining (SAM) Project, *Proc. 25th Symp. Geo-Environments and Geo-Techniques*, 113-118.


Code	MG0910-1	MG0910-3	MG0910-2
Run #	20373&20381	20374&20382	20375&20383
Sample	Sheep meat	White needle	Onion
Na	62 8.4	57 5.3	7286 249
Mg	14 5.2	5.4 3	1435 74
Si	9.2 3.8	4.6 2.3	ND
P	120 6.4	68 3.6	8341 265
S	406 17	92 5.1	2725 124
Cl	312 13	595 20	36836 1133
K	117 5.5	225 8.2	14740 458
Ca	30 3.1	33 2.1	4623 152
Ti	1.6 0.5	ND	ND
Fe	7.8 0.6	2.4 0.2	113 4.1
Cu	0.3 0.07	0.5 0.1	21 1.2
Zn	16 1.2	1.3 0.1	94 3.3
Br	0.9 0.2	1.3 0.1	61 2.8
Sr	ND	0.6 0.1	127 5.6

Table 1 PIXE analysis of food from Nariniiguur community, Bayan Obbo Soum*, Bayanhongor Aimag**, Mongolia Values in ppm, values in lower row indicate one sigma of the analysis. ND: not detected.


* A soum (Mongolian: сум) is the second level administrative subdivision below the aimags (provinces) in Mongolia.

** An aimag (аймаг) is an administrative subdivision in Mongolia, roughly comparable to Japan's prefecture. Mongolia is divided into 21 Aimags except for the capital Ulaanbaatar.

APPENDIX I



APPENDIX II



нар
дүн

1) Д. Байгальмаа, 37, эм, зах, 4-5 жилуу
800х
(88183532)
(88662207)

2) А. Уугатайеи, 18, эр, жилуу 2 жил
(88183532)
(88662207)

3) Н. Уугунбат, 64, эм, хирург, 7 жил
7 жил нувара, т02000
(98667927)
(89170771)